

Protocol 10-CBA

A Multicenter Access and Distribution Protocol for Unlicensed Cryopreserved Cord Blood Units (CBUs) for Transplantation in Pediatric and Adult Patients with Hematologic Malignancies and Other Indications

**Center for International Blood and Marrow Transplant Research (CIBMTR)
Clinical Research Organization (CRO) Services**

IND Number: 7555

NCT Number: NCT01351545

Version Number: v.10.5
25 Jul 2025

Sponsor Representatives
Heather Stefanski, MD, PhD

Sponsored by NMDPSM

Contact Information

Role	Contact Information
CIBMTR Protocol Officer	Heather Stefanski, MD, PhD NMDP/CIBMTR hstefans@nmdp.org (763) 406-8465 500 N 5th St. Minneapolis, MN 55401
Study Inbox	10-CBA Study Team NMDP/CIBMTR 10-CBA@nmdp.org
Study Web Pages	https://cibmtr.org/CIBMTR/Studies/Research-Programs/Clinical-Trials-Support/RCI-BMT/10-CBA-Study-Page https://network.nmdp.org/policies-protocols/cord-blood-banks/cord-blood-protocol-10-cba

Table of Contents

1	PROTOCOL SUMMARY	5
1.1	Protocol Synopsis	5
1.2	Schema.....	7
2	INTRODUCTION	8
2.1	Background.....	8
2.2	Rationale	8
3	STUDY DESIGN	9
4	STUDY OBJECTIVES.....	10
4.1	Primary Objective.....	10
4.2	Secondary Objectives	10
5	STUDY POPULATION.....	10
5.1	Inclusion Criteria.....	10
5.2	Exclusion Criteria	10
6	STUDY TREATMENT	11
6.1	Graft Selection.....	11
6.1.1	Graft Selection Inclusion Criteria	11
6.1.2	Graft Selection Exclusion Criteria.....	11
6.2	Dosage and Handling.....	12
6.3	Thaw and Infusion Treatment Plan	12
6.3.1	CBU Manipulation	12
6.3.2	CBU Administration.....	13
6.3.3	Multiple Cord Blood Transplants.....	13
6.3.4	Infusion Toxicities.....	13
6.4	Patient Conditioning, Transplant, and Post-Transplant Care.....	13
6.5	Toxicities and Risks	13
7	Patient Registration and Enrollment	14
8	Safety.....	14
8.1	Data Safety and Monitoring	14
8.2	Reporting Product Complaint Data.....	14
8.3	Reporting Serious Adverse Events	15
8.3.1	Reporting Instructions for the Investigator	15
8.4	NMDP Reporting to the FDA.....	15
9	Data Management	16
9.1	Data Reporting	16
9.2	Data Capture Methods	16
9.2.1	CIBMTR Outcomes Data (FormsNet3).....	16
9.3	Protocol Deviations.....	17
9.4	Study Monitoring	17
10	STATISTICAL CONSIDERATIONS.....	17
10.1	Study Design.....	17
10.2	Study Endpoints	17

10.3	Data Analysis	17
10.4	Disposition of Subjects	18
10.4.1	Off-Study Criteria	18
10.5	Abbreviations	19
10.6	Protocol Amendment History	20
11	REFERENCES	20

STATEMENT OF COMPLIANCE

The trial will be carried out in accordance with International Conference on Harmonisation Good Clinical Practice (ICH GCP) and the United States (US) Code of Federal Regulations (CFR) applicable to clinical studies (45 CFR Part 46, 21 CFR Part 50, 21 CFR Part 54, 21 CFR Part 56, 21 CFR Part 312, and/or 21 CFR Part 812).

The protocol, informed consent form(s), recruitment materials, and all subject materials will be submitted to the NMDP Institutional Review Board (IRB) for review and approval for sites relying on the NMDP IRB as their IRB of record. Documentation of local IRB submissions will be collected for those sites relying on local IRBs. IRB approval of both the protocol and the consent form must be obtained before any subject is enrolled. Any amendment to the protocol will require review and approval by the IRB before the changes are implemented to the study with a determination for re-consent for previously enrolled subjects.


1 PROTOCOL SUMMARY

1.1 PROTOCOL SYNOPSIS

Protocol Title (Full)	A Multicenter Access and Distribution Protocol for Unlicensed Cryopreserved Cord Blood Units (CBUs) for Transplantation in Pediatric and Adult Patients with Hematologic Malignancies and Other Indications
Protocol Title (Short)	10-CBA
Study Design	This study is an access and distribution protocol for unlicensed CBUs in pediatric and adult patients with hematologic malignancies and other indications
Study Objective(s)	<p>Primary Objective: To examine the incidence of neutrophil recovery of $\geq 500/\text{mm}^3$ after cord blood transplantation in a multi-institution setting using CBUs that are not Food and Drug Administration (FDA) licensed</p> <p>Secondary Objectives:</p> <ul style="list-style-type: none"> • Assess incidence of transmission of infection • Assess incidence of serious infusion reaction • Determine overall survival 1 year after cord blood transplantation • Assess cumulative incidence of acute graft vs. host disease (GVHD) grades II to IV and grades III to IV • Assess cumulative incidence of chronic GVHD

	<ul style="list-style-type: none"> Determine platelet engraftment of >20,000 μL and >50,000 μL
Study Population	Pediatric and adult patients with hematologic malignancies and other indications
Treatment Description	Treatment, including pre-transplant conditioning and GVHD prophylaxis, will occur per each transplant center's (TC) specifications
Accrual Objective	In this access and distribution protocol, US patients undergoing transplantation with unlicensed CBUs will be enrolled and there is no accrual maximum
Principal Investigators	The principal investigators (PIs) are transplant physicians at all participating US TCs
Participation Duration	Subjects will be followed through 1 year after transplant
Accrual Period	The accrual period is open ended
Inclusion Criteria	<ol style="list-style-type: none"> Disorders affecting the hematopoietic system that are inherited, acquired, or result from myeloablative treatment Signed informed consent (and signed assent, if applicable) obtained prior to study enrollment Pediatric and adult patients of any age
Exclusion Criteria	<ol style="list-style-type: none"> Patients who are receiving only licensed CBUs Cord blood transplant recipients at international TCs Patients who are enrolled on another IND protocol to access each unlicensed CBU Patients whose selected unlicensed CBU(s) will be more than minimally manipulated
Statistical Methods	Descriptive analyses will be performed for primary transplants annually, as part of regular FDA IND annual reporting

1.2 SCHEMA



2 INTRODUCTION

2.1 BACKGROUND

Hematopoietic stem cell transplantation (HSCT) is now recognized as an effective form of therapy for an increasing number of malignant and non-malignant disorders.¹ Human umbilical cord blood (UCB) is an alternative source of hematopoietic stem cells (HSCs) that is capable of reconstituting hematopoiesis after intensive myeloablative therapy, thereby extending the unrelated donor pool.²⁻¹⁰

Cell dose is the most important factor affecting clinical outcomes in adult patients undergoing transplantation with a single CBU.¹¹⁻¹⁴ Transplantation with a cell dose of less than 2.5×10^7 nucleated cells per kilogram or a CD34 dose of less than 1.7×10^5 per kilogram is associated with higher rates of non-engraftment, non-relapse mortality (NRM) and lower survival.¹⁴ In pediatric patients, 5-year leukemia free survival following transplantation of CBU(s) with a single or double human leukocyte antigen (HLA) mismatch was comparable to transplantation with matched or single HLA-mismatched marrow and survival was possibly higher with fully HLA-matched cord blood.¹⁵ In adult patients, transplantation with CBU(s) with up to 2 HLA mismatches had comparable leukemia-free survival to transplantation with matched or single HLA-mismatched unrelated peripheral blood stem cells (PBSC) and marrow.¹⁶ However, in patients receiving cord blood, transplant-related mortality (TRM) was higher and grades 2-4 acute GVHD was lower than in patients receiving fully matched PBSC or marrow.

UCB has advantages when compared to grafts from adult cell sources including being rapidly available, having low rates of viral contamination, and having lower probabilities of severe acute and extensive chronic GVHD despite the use of HLA-mismatched grafts. However, a major disadvantage is that low graft cell dose in adult recipients leads to delayed hematopoietic recovery, an increased risk of graft failure and increased risk for TRM, which limits the application of UCB transplant in adults. Barker and colleagues at The University of Minnesota has investigated the novel approach of the combined transplantation of two CBUs in a double unit graft after myeloablative conditioning as a strategy to augment graft cell dose.¹⁷ In this study, all 21 patients receiving two CBUs engrafted at a median of 21 days and while both CBUs engrafted initially, by Day 100 one CBU predominated. Subsequent studies have demonstrated decreased relapse, increased NRM and similar leukemia-free survival in recipients of double cord transplants when compared to patients receiving matched unrelated or matched related PBSC and marrow.¹⁸⁻²⁰

These and other studies demonstrate that UCB is safe and effective for transplantation. Based on this clinical evidence, the FDA has released criteria that will allow CBUs that meet the criteria to be distributed as a licensed biologic drug.

2.2 RATIONALE

In 1997, the FDA announced a risk based approach to the regulation of human cellular and tissue-based products including UCB.^{21,22} The approach was finalized as the three final rules for Human Cells, Tissues, and Tissue-Based Products (HCT/Ps) that were effective on March 29, 2004 (requirement for establishment registration) and on May 25, 2005 (donor eligibility and Good Tissue Practices).²³⁻²⁵ These rules were designed to: prevent the transmission of communicable disease, minimize contamination and

preserve integrity and function during processing, outline safety and effectiveness requirements for cells from unrelated donors or when HCT/PS are more than minimally manipulated, assure labeling is clear, accurate and not misleading and monitor and communicate with industry via establishment registration.²²

In October 2009, the FDA released a final licensure guidance entitled *Minimally Manipulated, Unrelated Allogeneic Placental/Umbilical Cord Blood Intended for Hematopoietic Reconstitution for Specified Indications*, which outlines ways a cord blood bank (CBB) may apply for licensure of unrelated, allogeneic CBU.²⁶ CBBs may submit a biologics license application (BLA) to manufacture licensed CBUs as defined in the guidance for specified hematologic and non-hematologic disorders. CBUs meeting the criteria will be considered licensed and will be able to be distributed as a licensed biological drug.

However, as of October 20, 2011, those CBUs that do not meet the manufacturing requirements for licensure can only be distributed for transplantation if the transplant will occur under an IND research protocol. These CBUs are in current and future inventory at domestic and international CBBs that can't be demonstrated to meet licensing requirements.

In addition to the licensure guidance, the FDA published a companion guidance in August 2011 entitled *Investigational New Drug Applications (INDs) for Minimally Manipulated, Unrelated Allogeneic Placental/Umbilical Cord Blood Intended for Hematopoietic Reconstitution for Specified Indication*.²⁷ This guidance provides advice to assist sponsors in submission of an IND application for distribution of CBUs that do not meet licensure requirements when they are needed for transplantation of a patient with a serious or life-threatening disease and there is no satisfactory alternative treatment. Sponsors of an IND may include CBBs, registries (such as NMDP), and individual physicians serving as sponsor-investigators.

After licensing criteria are implemented, there will be an extensive current inventory of CBUs that does not meet the licensing criteria but has a high likelihood of containing the CBU that is the best available match for a patient. Accordingly, the purpose of this protocol is to provide access to, and distribution of CBUs that do not meet the manufacturing requirements for licensure when they are the best available option for a patient with a serious or life-threatening disease in need of transplantation. This IND provides a single, efficient model that benefits from NMDP established relationships with an extensive network of CBBs, registries, TCs and physicians, yet it does not limit options for other potential IND sponsors. The protocol will allow both US and international CBBs to distribute CBUs under this IND as long as they meet the requirements established by the NMDP. It is likely that the majority of CBUs collected previously will not meet licensure requirements and will need to be distributed under this (or another) IND. Similarly, while approximately 17% of domestic transplants involved a CBU from an international CBB in 2009, it is anticipated that most of these CBBs will chose not to hold an IND for distribution of a small number of CBUs to the US.

3 STUDY DESIGN

This is an access and distribution protocol for unlicensed cryopreserved CBUs for transplant of pediatric and adult patients with hematologic malignancies and other indications. The study will enroll an unlimited number of transplant recipients at participating TCs based on well-defined inclusion and exclusion criteria, when the best graft source is determined to be a domestic or international CBU that

does not meet FDA licensure requirements. The follow-up schedule and affiliated data collection forms for the unlicensed cord blood recipients are the standard CIBMTR data forms and procedures.

All recipients receiving at least one unlicensed CBU facilitated under the NMDP IND will be enrolled in the study and will be assessed for: neutrophil engraftment, serious infusion reaction, transmission of infection from CBU, survival, acute and chronic GVHD, and platelet engraftment.

4 STUDY OBJECTIVES

4.1 PRIMARY OBJECTIVE

The primary objective of this study is to examine the incidence of neutrophil recovery of $\geq 500/\text{mm}^3$ after cord blood transplantation in a multi-institution setting using CBUs that are not FDA licensed.

4.2 SECONDARY OBJECTIVES

The secondary objectives are as follows:

- Assess incidence of transmission of infection
- Assess incidence of serious infusion reaction
- Determine overall survival 1 year after cord blood transplantation
- Assess cumulative incidence of acute GVHD grades II to IV and grades III to IV
- Assess cumulative incidence of chronic GVHD
- Determine platelet engraftment of $>20,000 \mu\text{L}$ and $>50,000 \mu\text{L}$

5 STUDY POPULATION

5.1 INCLUSION CRITERIA

To be eligible to participate in this study, an individual must meet all the following criteria:

1. Disorders affecting the hematopoietic system that are inherited, acquired, or result from myeloablative treatment
2. Signed informed consent (and signed assent, if applicable) obtained prior to study enrollment
3. Pediatric and adult patients of any age

5.2 EXCLUSION CRITERIA

An individual who meets any of the following criteria will be excluded from participation in this study:

1. Patients who are receiving only licensed CBUs
2. Cord blood transplant recipients at international TCs
3. Patients who are enrolled on another IND protocol to access each unlicensed CBU
4. Patients whose selected unlicensed CBU(s) will be more than minimally manipulated

6 STUDY TREATMENT

6.1 GRAFT SELECTION

Criteria for the CBU(s) selected will be determined by the TC. The selected CBU, or at least one of the selected CBUs for a multiple cord blood transplant, must be unlicensed.

6.1.1 GRAFT SELECTION INCLUSION CRITERIA

Unlicensed CBUs selected by the TC must include:

1. For a single CBU transplant, a post-processing dose of $\geq 2.5 \times 10^7$ total nucleated cells (TNC)/kg recipient weight is recommended, with minimum allowable dose of $\geq 1.0 \times 10^7$ TNC/kg recipient weight
2. Proof of sterility – sterility will be reported at time of release
3. Results of an assay for cell viability (e.g., colony forming units (CFU), CD34+ viability, Trypan Blue or other staining method(s) or another functional assay supporting cell viability)
 - a. If a post-processing / pre-freeze cell viability assay was not performed by the CBB or does not meet current minimum requirements, cell viability from a thawed retention sample or segment representative of the cord blood product will be performed prior to transplantation
 - b. If no cell viability assay can be performed, the TC will confirm the CBU remains the best match for the patient for transplant
4. Any hemoglobinopathy testing done by the CBB (including testing for sickle cell and beta thalassemia) will be communicated with the TC
5. Identity testing: HLA-A, -B, -C, -DRB1 by DNA-based technologies at a resolution sufficient to determine concordance with the original HLA typing must be confirmed prior to transplant
 - a. High resolution HLA-DRB1 typing must be performed prior to shipment of the CBU and may be performed at the recruitment, extended typing or confirmatory/verification typing stages
 - b. This testing may be performed in a laboratory designated by the CBB or NMDP or another national registry
 - c. The laboratory performing this testing must be capable of carrying out DNA-based HLA typing and be accredited by ASHI, EFI, or CAP
6. It is recommended that each CBU is at least 4/6 HLA-matched at HLA-A, -B, and -DRB1, with a minimum allowable HLA match of 3/6 at HLA-A, -B, and -DRB1

6.1.2 GRAFT SELECTION EXCLUSION CRITERIA

Unlicensed CBUs selected by the TC must not include:

1. Compromised traceability to product or maternal identity suggestive of mix-up such as discrepant HLA typing on identity confirmation
2. Positive microbial culture of CBU (bacterial and/or fungal)
3. Positive hemoglobinopathy testing indicating CBU is:
 - a. Homozygous for sickle cell and/or β thalassemia disease

- b. Heterozygous for both sickle cell and β thalassemia trait
 - c. Severe alpha thalassemia (hemoglobin H disease)
- 4. Positive/reactive result for Infectious Disease Markers (IDM) (FDA licensed/cleared/approved donor screening test or other laboratory IDM test as part of relevant medical records):
 - a. Human immunodeficiency virus (HIV) types 1 and 2
 - b. Hepatitis B Virus (HBV)
 - i. Hepatitis B Surface Antigen (HBsAg) test
 - ii. HBV NAT test
 - c. Hepatitis C Virus (HCV)
 - d. T cruzi/Chagas Disease
 - e. West Nile Virus (WNV)
- 5. No Infectious disease test result for:
 - a. HIV type 1 /type 2 antibody test
 - b. HIV p24 Antigen or HIV nucleic acid test (NAT)
 - c. HBsAg test
 - d. HCV antibody test
- 6. No bacterial sterility testing performed on CBU
- 7. Collection and storage of CBU(s) in vials

6.2 DOSAGE AND HANDLING

For a single CBU transplant, a post-processing dose of $\geq 2.5 \times 10^7$ TNC/kg recipient weight is recommended, with minimum allowable dose of $\geq 1.0 \times 10^7$ TNC/kg recipient weight. The maximum dose of dimethyl sulfoxide (DMSO) should not exceed 1mL (gram) of DMSO per kilogram of recipient weight per day of administration.

The CBU should be handled, stored, and prepared according to the manufacturer's instructions or validated institution procedures. In the case of transplantation of multiple CBUs, the subsequent CBU(s) should not be thawed until it is confirmed that the patient has recovered from the side effects from the prior CBU infused.

Supplies and reagents that come in contact with the cellular therapy product shall be of the appropriate quality/grade for the intended use and, whenever possible, be FDA-approved. Supplies and reagents that are not FDA approved, but are available outside the US, such as Dextran in normal saline, may be used under this protocol.

6.3 THAW AND INFUSION TREATMENT PLAN

Recommendations for preparation and infusion of CBU are outlined below.

6.3.1 CBU MANIPULATION

Minimal manipulation is allowed and should be conducted per the manufacturer's instructions or validated institution procedures.

- Minimal manipulation is defined by current FDA guidelines

- CBUs that are not red cell reduced must be washed
- Thaw and reconstitution or wash methods should be based on a sufficient dilution ratio based on the original cord blood volume

6.3.1.1 PREMEDICATIONS

The patient should be premedicated prior to infusion per institution standard operating procedures.

6.3.2 CBU ADMINISTRATION

The CBU intravenous infusion should be performed according to manufacturer's instructions or per institution standard operating procedures.

- A standard blood filter should be used
- The time from the initiation of thaw to completion of infusion should be minimized, preferably within 2 hours

6.3.3 MULTIPLE CORD BLOOD TRANSPLANTS

- Thaw CBUs independently, thawing the second CBU after the first CBU is successfully infused
- Infuse CBUs independently
- Should a reaction occur resulting from infusion of the first CBU, appropriately manage the reaction before second CBU is thawed for infusion

6.3.4 INFUSION TOXICITIES

Patient monitoring is recommended during the infusion, per institution standard operating procedures.

The infusion should be discontinued in the event of serious toxicity and/or product complaint. The infusion is discontinued based on clinician assessment and in accordance with institution standard operating procedures. See Section 8 for adverse event (AE) and product complaint reporting details.

6.4 PATIENT CONDITIONING, TRANSPLANT, AND POST-TRANSPLANT CARE

The conditioning regimen, transplant, GVHD prophylaxis and supportive care for the transplant patient will be determined by the individual institution performing the transplant and reported to the CIBMTR on the appropriate case report forms (CRFs).

6.5 TOXICITIES AND RISKS

Transplant-related toxicities attributed to the product infusion will be reported to the CIBMTR on the appropriate CRFs. Toxicities potentially associated with the infusion of the CBU include DMSO toxicity and side effects from intact and hemolyzed red cells and may include changes in heart rate or rhythm, changes in blood pressure, fever, chills, sweats, nausea/vomiting, diarrhea, abdominal cramping, fluid overload, headache, dyspnea, hemoglobinuria, allergic reaction, acute renal failure, and in rare cases

infusion reaction resulting in death. If toxicities are serious, they must be reported in an expedited fashion to NMDP per the reporting instructions outlined in Section 8.3.1.

Transplant-related risks attributed to receiving CBUs include delayed neutrophil and platelet recovery, and lower rates of engraftment compared with PBSC or marrow. In addition, cord blood is a single-source product; industry-standard practices assure donor mothers that their children will never be contacted for an additional donation request. Cord blood transplant complications that could be best managed with additional nucleated cells require novel strategies such as additional CBUs or a switch to an adult donor. Rarer risks can include seroconversion or development of a communicable disease (limited by stringent IDM screening), or transmission of a rare disease or genetic disorder to the recipient due to scant medical history on the newborn donor.

7 PATIENT REGISTRATION AND ENROLLMENT

In instances where an unlicensed CBU is determined to be the best transplant option, patient enrollment in this study is required when the selected unlicensed CBU is facilitated under the NMDP IND. Facilitation includes NMDP facilitated the procurement, collection, and/or transportation of the product.

All participants in this access and distribution protocol must provide informed consent on the Informed Consent Form (ICF) approved by the IRB of record and the 10-CBA study team, prior to study enrollment. At the time of CBU order, the TC will enroll the patient in line with Section 9.

If a previously enrolled participant requires a subsequent transplant using unlicensed CBUs and at least one CBU is procured under the NMDP IND, the patient must be re-consented and re-enrolled in the study at the time of CBU order.

8 SAFETY

8.1 DATA SAFETY AND MONITORING

The NMDP Donor and Patient Safety Monitoring Advisory Group (DPSM) will review a summary of adverse events regularly. The report will be made available to participating TCs. If the DPSM recommends protocol or informed consent changes, they will be distributed to the participating PIs. It is the responsibility of each PI to forward the distributed communication to their local Institutional Review Board (IRB) per local reporting policies.

8.2 REPORTING PRODUCT COMPLAINT DATA

Upon receipt of the CBU, the TC will complete written confirmation that the product was received. If the integrity of the CBU is compromised at receipt (or at any time after receipt), the TC should report to NMDP within 3 business days using the Product Complaint form in the FormsNet3 Recipient module.

Examples include: a broken CBU, compromised shipping conditions, a contaminated product, or viability lower than expected. NMDP will notify the CBB and conduct an investigation per NMDP standard

procedures for these complaints. NMDP will report manufacturing complaints that are determined to be related to the manufacturing of a distributed CBU to the FDA annually.

NMDP as trial sponsor and central monitoring entity will be responsible for reporting all adverse events that meet Office of Human Research Protections (OHRP) definition of unanticipated problem to OHRP.

8.3 REPORTING SERIOUS ADVERSE EVENTS

Because all or most US recipients participating on this protocol will be receiving potentially toxic preparative therapy, significant regimen-related toxicity is anticipated. Likewise, substantial mortality and non-engraftment of a transplanted CBU are anticipated. This data will be captured via filing of appropriate CIBMTR CRFs. CIBMTR will not report non-engraftment or complications that are attributed to the natural course of the disease, relapse, or chemotherapy and/or radiation toxicity.

8.3.1 REPORTING INSTRUCTIONS FOR THE INVESTIGATOR

All serious adverse events (SAEs) categorized below will be reported in accordance with the FDA requirements for AE reporting.

All SAEs determined to be caused by or probably caused by the CBU based on objective evidence must be reported promptly under this IND protocol using the Adverse Event form in the FormsNet3 Recipient module.

Reporting of SAEs that result in death or a life-threatening event should occur as soon as possible, but no later than 24 hours of learning of the event. Other SAEs should be reported within 3 working days.

- Recipient seroconversion to any of the FDA-listed relevant communicable diseases within 6 months of CBU infusion which, upon investigation, is determined to be caused or potentially caused by the CBU
- Recipient bacteremia related to a contaminated CBU
- Recipient develops any of the FDA-listed relevant communicable diseases within 6 months of CBU infusion which, upon investigation, is determined to be caused or potentially caused by the CBU
- Serious infusion reaction within first 24 hours after infusion

8.4 NMDP REPORTING TO THE FDA

A medical monitor will review all SAEs reported by the clinical trial site and be responsible for determining severity and relationship to the product.

Any SAE that is determined to be possibly linked to the product will be investigated by NMDP Quality Assurance Nurses to determine if there is a causal relationship requiring expedited reporting. NMDP will submit expedited reporting to FDA only for unexpected SAEs and for expected SAEs that occur at a frequency higher than usual.

If, during the investigation of the SAE, a problem(s) with manufacturing practices is discovered, NMDP will contact the appropriate CBB to initiate corrective action for the CBB-specific procedures and report under 8.2 above, as applicable.

9 DATA MANAGEMENT

9.1 DATA REPORTING

The data collection forms for the subjects enrolled on this study include the standard CIBMTR data collection forms (including Adverse Events and Product Complaints meeting study reporting requirements) in the FormsNet3 Recipient module, and study specific CIBMTR data collection forms in the Electronic Data Capture (EDC) application.

Many important data elements for the study are collected on the standard CIBMTR reporting forms, and therefore timely and accurate completion of these forms is essential.

9.2 DATA CAPTURE METHODS

9.2.1 CIBMTR OUTCOMES DATA (FORMSNET3)

All TCs participating in this protocol are already reporting baseline and outcome data on all transplant recipients to CIBMTR. Data reporting requirements for this protocol will be fulfilled by those data. There are no additional follow-up data requirements for this protocol.

CIBMTR data reporting time points used in this study:

- Baseline (at time of transplant)
- Infusion data (at time of transplant)
- 100 days post-transplant
- 6 months post-transplant
- 1-year post-transplant
- At time of patient's death

TCs must continue to complete standard CIBMTR follow-up reporting on these patients beyond the study time point of 1-year per CIBMTR reporting requirements.

9.2.1.1 FORMSNET3 CIBMTR RECIPIENT FORMS

Please refer to the Forms Instructions Manual for required forms.

Standard CIBMTR data report forms can be found at: <https://cibmtr.org/CIBMTR/Data-Operations/Data-Collection-Forms>

9.2.1.2 EDC FORMS

Please refer to the CRF Complete Guidelines (CCGs) for instructions on how to complete the study-specific forms in the EDC system.

9.3 PROTOCOL DEVIATIONS

When an emergency occurs that requires a deviation from the protocol for patient safety, a decision will be made as soon as possible to determine whether or not the subject (for whom the deviation from protocol was affected) is to continue in the study. All protocol deviations (PDs) will be documented in the EDC System.

Study sites are also required to note deviations in patient medical records and report them to their local IRB in accordance with local policies. All protocol deviations will be compiled centrally by the CIBMTR Protocol Coordinator and reported to the NMDP DPSM.

Planned Protocol deviations should be discussed with the 10-CBA study team via email at 10-CBA@nmdp.org.

9.4 STUDY MONITORING

The PI will permit study-related remote and/or centralized monitoring visits by representatives of the CIBMTR or designees, and regulatory inspection(s) (e.g., FDA) to ensure proper conduct of the study and compliance with all FDA safety reporting requirements. Access must be provided to source documents, data collection forms, consent and assent forms, and any other study documents.

10 STATISTICAL CONSIDERATIONS

10.1 STUDY DESIGN

This study is a multicenter access and distribution protocol, to allow access to cord blood transplantation for patients whose best graft source is determined to be a CBU from US or international CBBs not meeting FDA licensure requirements. Patients will be enrolled and followed prospectively to assess outcomes after transplant, from data collected using CIBMTR forms. The study is observational since the choice of the best available CBU will be left to the discretion of the patient's transplant physician. As the registry of licensed CBU grows, enrollment on this protocol is expected to gradually decline in relation to the overall number of CBUs used for transplantation. This study is descriptive in nature and there is no accrual limit.

10.2 STUDY ENDPOINTS

The primary endpoint is neutrophil engraftment. Other important secondary endpoints include platelet engraftment, acute GVHD grades II-IV or III-IV, chronic GVHD, overall survival at 1 year, transmission of infection, and serious infusion reaction.

10.3 DATA ANALYSIS

Interim descriptive analyses will be performed for primary transplants annually, as part of regular IND annual reporting. As the eligibility for this protocol is very broad, analyses will be done separately for several pre-specified subgroups. First, analyses will be conducted separately for patients transplanted for malignant diseases and those transplanted for non-malignant diseases. They will also be analyzed separately for transplants using single CBU and those using multiple CBUs. Further separation of the study population such as adults vs. pediatrics and myeloablative vs. nonmyeloablative conditioning regimens will be considered depending on the patient numbers. In particular, neutrophil and platelet engraftment will only be analyzed using transplants with myeloablative conditioning regimens.

Baseline characteristics will be described using frequencies/percents or median/range as appropriate.

Transplant outcomes will be described with confidence intervals as follows. Probability of neutrophil and platelet engraftment, and probability of developing acute or chronic GVHD will be described using the cumulative incidence estimate²⁸, with death prior to engraftment or death prior to development of GVHD as competing risks, respectively. Overall survival will be described using the Kaplan-Meier estimator.²⁹ In addition, a descriptive summary of product related relevant communicable disease agents or diseases (RCDAD) will be provided as well as listing of patients who experience serious infusion toxicity within 24 hours.

Due to the fact that the previous NMDP CBU protocol, *A centralized cord blood registry to facilitate allogeneic unrelated donor umbilical cord blood transplantation*, contained stopping rules, many years of data on the safety and efficacy of umbilical cord transplant using CBUs in our inventory have been reported. These data have shown that umbilical cord transplant is a comparably safe and effective alternative to bone marrow and PBSC transplantation. Therefore, stopping rules are not necessary for this protocol.

10.4 DISPOSITION OF SUBJECTS

The number of subjects enrolled in the study and the disposition of all subjects will be summarized. Subjects who discontinue study will be listed according to the off-study criterion that applies.

10.4.1 OFF-STUDY CRITERIA

Study sites must complete the Study Exit form in the Medidata Rave application for any subjects meeting one of these criteria:

1. Subject withdrawal
2. Transplant canceled: CBU(s) **not** shipped under NMDP IND
3. Substituted non-NMDP IND CBU: CBU(s) **not** shipped under NMDP IND
4. Substituted other cell source (BM, PBSC): CBU(s) **not** shipped under NMDP IND
5. Subject determined ineligible

Study subjects should **not** be exited after CBU(s) shipped under the NMDP IND. Subjects should **not** be exited for the following reasons:

1. Subject death *after* CBU(s) shipped under the NMDP IND
2. Transplant canceled *after* CBU(s) shipped under the NMDP IND

3. Substituted non-NMDP IND CBU(s) *after* CBU(s) shipped under the NMDP IND
4. Substituted other cell source (BM, PBSC) *after* CBU(s) shipped under the NMDP IND

Please contact the 10-CBA study team at 10-CBA@nmdp.org with any questions or concerns regarding study exits.

10.5 ABBREVIATIONS

AE	Adverse Event
BLA	Biologics License Application
CBB	Cord Blood Bank
CBU	Cord Blood Unit
CFR	Code of Federal Regulations
CFU	Colony Forming Units
CIBMTR	Center for International Blood and Marrow Transplant Research
CRF	Case Report Form
CRO	Clinical Research Organization
DMSO	Dimethyl Sulfoxide
DPSM	Donor and Patient Safety Monitoring Advisory Group
EDC	Electronic Data Capture
FDA	Food and Drug Administration
GCP	Good Clinical Practice
GVHD	Graft Versus Host Disease
HBsAg	Hepatitis B Surface Antigen
HCT/P	Human Cells, Tissues, and Tissue-Based Product
HCV	Hepatitis C Virus
HIV	Human Immunodeficiency Virus
HLA	Human Leukocyte Antigen
HSC	Hematopoietic Stem Cell
HSCT	Hematopoietic Stem Cell Transplantation
ICH	International Conference on Harmonisation
IDM	Infectious Disease Markers
IND	Investigational New Drug
IRB	Institutional Review Board
NAT	Nucleic Acid Test
NCT	National Clinical Trial
NRM	Non-Relapse Mortality
OHRP	Office of Human Research Protections
PBSC	Peripheral Blood Stem Cell

PD	Protocol Deviation
PI	Principal Investigator
RCDAD	Relevant Communicable Disease Agents or Diseases
SAE	Serious Adverse Event
TC	Transplant Center
TNC	Total Nucleated Cells
TRM	Transplant-Related Mortality
UCB	Umbilical Cord Blood
US	United States
WNV	West Nile Virus

10.6 PROTOCOL AMENDMENT HISTORY

Version	Date	Description of Change	Brief Rationale
10.5	25 JUL 2025	Protocol template update. Removed Dr. Miller's name and contact info for Dr. Stefanski's contact info as the new study-level PI. Removed Medidata Rave EDC language and CRF reporting table. Added statement of compliance language, corrected protocol coordinator to 10-CBA study team and corrected spelling or grammatical errors.	The updates for this amendment were made primarily because of a study level PI change and an EDC system change from Medidata Rave to Veeva CDMS. The formatting changes were due to updating the protocol template to reflect current company quality standards.

11 REFERENCES

¹Appelbaum FR, Forman SJ, Negrin RS, Blume KG. Thomas' Hematopoietic Cell Transplantation. 4 ed: Wiley-Blackwell; 2009.

²Kurtzberg J, Laughlin M, Graham ML, Smith C, Olson JF, Halperin EC, et al. Placental blood as a source of hematopoietic stem cells for transplantation into unrelated recipients. N Engl J Med. 1996 Jul 18;335(3):157-66.

³Wagner JE, Rosenthal J, Sweetman R, Shu XO, Davies SM, Ramsay NK, et al. Successful transplantation of HLA-matched and HLA-mismatched umbilical cord blood from unrelated donors: analysis of engraftment and acute graft-versus-host disease. Blood. 1996 Aug 1;88(3):795-802.

⁴Laughlin MJ, Barker J, Bambach B, Koc ON, Rizzieri DA, Wagner JE, et al. Hematopoietic engraftment

- and survival in adult recipients of umbilical-cord blood from unrelated donors. *N Engl J Med*. 2001 Jun 14;344(24):1815-22.
- ⁵Laughlin MJ, Eapen M, Rubinstein P, Wagner JE, Zhang MJ, Champlin RE, et al. Outcomes after transplantation of cord blood or bone marrow from unrelated donors in adults with leukemia. *N Engl J Med*. 2004 Nov 25;351(22):2265-75.
- ⁶Rocha V, Labopin M, Sanz G, Arcese W, Schwerdtfeger R, Bosi A, et al. Transplants of umbilical-cord blood or bone marrow from unrelated donors in adults with acute leukemia. *N Engl J Med*. 2004 Nov 25;351(22):2276-85.
- ⁷Takahashi S, Iseki T, Ooi J, Tomonari A, Takasugi K, Shimohakamada Y, et al. Single-institute comparative analysis of unrelated bone marrow transplantation and cord blood transplantation for adult patients with hematologic malignancies. *Blood*. 2004 Dec 1;104(12):3813-20.
- ⁸Barker JN, Weisdorf DJ, Defor TE, Blazar BR, Miller JS WJ. Rapid and complete donor chimerism in adult recipients of unrelated donor umbilical cord blood transplantation after reduced-intensity conditioning. *Blood* 2003;102:1915-9.
- ⁹Takahashi S, Ooi J, Tomonari A, Konuma T, Tsukada N, Oiwa-Monna M, et al. Comparative single-institute analysis of cord blood transplantation from unrelated donors with bone marrow or peripheral blood stem-cell transplants from related donors in adult patients with hematologic malignancies after myeloablative conditioning regimen. *Blood*. 2007 Feb 1;109(3):1322-30.
- ¹⁰Atsuta Y, Suzuki R, Nagamura-Inoue T, Taniguchi S, Takahashi S, Kai S, et al. Disease-specific analyses of unrelated cord blood transplantation compared with unrelated bone marrow transplantation in adult patients with acute leukemia. *Blood*. 2009 Feb 19;113(8):1631-8.
- ¹¹Gluckman E, Rocha V, Boyer-Chamard A, Locatelli F, Arcese W, Pasquini R, et al. Outcome of cord-blood transplantation from related and unrelated donors. Eurocord Transplant Group and the European Blood and Marrow Transplantation Group. *N Engl J Med*. 1997 Aug 7;337(6):373-81.
- ¹²Rubinstein P, Carrier C, Scaradavou A, Kurtzberg J, Adamson J, Migliaccio AR, et al. Outcomes among 562 recipients of placental-blood transplants from unrelated donors. *N Engl J Med*. 1998 Nov 26;339(22):1565-77.
- ¹³Wagner JE, Barker JN, DeFor TE, Baker KS, Blazar BR, Eide C, et al. Transplantation of unrelated donor umbilical cord blood in 102 patients with malignant and nonmalignant diseases: influence of CD34 cell dose and HLA disparity on treatment-related mortality and survival. *Blood*. 2002 Sep 1;100(5):1611-8.
- ¹⁴Rocha V, Gluckman E. Improving outcomes of cord blood transplantation: HLA matching, cell dose and other graft- and transplantation-related factors. *Br J Haematol*. 2009 Oct;147(2):262-74.
- ¹⁵Eapen M, Rubinstein P, Zhang MJ, Stevens C, Kurtzberg J, Scaradavou A, et al. Outcomes of

transplantation of unrelated donor umbilical cord blood and bone marrow in children with acute leukaemia: a comparison study. *Lancet*. 2007 Jun 9;369(9577):1947-54.

- ¹⁶Eapen M, Rocha V, Sanz G, Scaradavou A, Zhang MJ, Arcese W, et al. Effect of graft source on unrelated donor haemopoietic stem-cell transplantation in adults with acute leukaemia: a retrospective analysis. *Lancet Oncol*. 2010 Jul;11(7):653-60.
- ¹⁷Barker JN, Weisdorf DJ, DeFor TE, Blazar BR, McGlave PB, Miller JS, et al. Transplantation of 2 partially HLA-matched umbilical cord blood units to enhance engraftment in adults with hematologic malignancy. *Blood*. 2005 Feb 1;105(3):1343-7.
- ¹⁸Gutman JA, Leisenring W, Appelbaum FR, Woolfrey AE, Delaney C. Low relapse without excessive transplant-related mortality following myeloablative cord blood transplantation for acute leukemia in complete remission: a matched cohort analysis. *Biol Blood Marrow Transplant*. 2009 Sep;15(9):1122-9.
- ¹⁹Brunstein CG, Gutman JA, Weisdorf DJ, Woolfrey AE, DeFor TE, Gooley TA, et al. Allogeneic hematopoietic cell transplantation for hematologic malignancy: relative risks and benefits of double umbilical cord blood. *Blood*. 2010 Nov 25;116(22):4693-9.
- ²⁰Ponce DM, Zheng J, Gonzales A, Lubin M, Heller G, Castro-Malaspina H, et al. Reduced Late Mortality Risk Contributes to Similar Survival after Double Unit Cord Blood Transplantation as Compared with Related and Unrelated Donor Hematopoietic Stem Cell Transplantation. *Biol Blood Marrow Transplant*. Jan 10.
- ²¹Food and Drug Administration. Proposed Approach to Regulation of Cellular and Tissue-Based Products (docket number 97N-0068). Rockville (MD): FDA.gov [Internet]; 1997.
<http://www.fda.gov/downloads/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/Rissue/UCM062601.pdf>
- ²²Havarth L. A Brief History of FDA Regulation of Human Cells and Tissues. In: Areman E, Loper K, editors. *Cellular Therapy: Principles, Methods, and Regulations*. Bethesda: AABB; 2009.
- ²³Food and Drug Administration. Human Cells, Tissues, and Cellular and Tissue-Based Products; Establishment Registration and Listing (interim final rule) (69 FR 3823, January 27, 2004).
- ²⁴Food and Drug Administration. Eligibility Determination for Donors of Human Cells, Tissues, and Cellular and Tissue-Based Products (donor eligibility final rule) (69 FR 29786, May 25, 2004).
- ²⁵Food and Drug Administration. Current Good Tissue Practice for Human Cell, Tissue, and Cellular and Tissue-Based Product Establishments, Inspection and Enforcement (CGTP final rule) (69 FR 68612, November 24, 2004).
- ²⁶Food and Drug Administration. Guidance for Industry: Minimally Manipulated, Unrelated Allogeneic

Placental/Umbilical Cord Blood Intended for Hematopoietic Reconstitution for Specified Indications (October 2009).

<http://www.fda.gov/downloads/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/Blood/UCM187144.pdf>

²⁷Food and Drug Administration. Guidance for Industry and FDA Staff: Investigational New Drug Applications (INDs) for Minimally Manipulated, Unrelated Allogeneic Placental/Umbilical Cord Blood Intended for Hematopoietic Reconstitution for Specified Indications (June 2011).
<http://www.fda.gov/downloads/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/Blood/UCM187146.pdf>

²⁸Gooley TA, Leisenring W, Crowley J, Storer BE. Estimation of failure probabilities in the presence of competing risks: new representations of old estimators. Stat Med. 1999 Mar 30;18(6):695-706.

²⁹Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. J Am Stat Assoc. 1958;53:695-706.